

# THE STRUCTURE AND STAR FORMATION HISTORY OF EARLY-TYPE GALAXIES IN THE UDF/GRAPES SURVEY

A. PASQUALI

Institut für Astronomie, HPF, ETH, 8093 Zürich, Switzerland

I. FERRERAS

Department of Physics and Astronomy, University College London, Gower St. London WC1E 6BT, England

N. PANAGIA, S. MALHOTRA, J.E. RHOADS, N. PIRZKAL, AND A.M. KOEKEMOER, L. MOUSTAKAS, C. XU  
STScI, 3700 San Martin Drive, Baltimore, MD 21218, USA

E. DADDI

Spitzer Fellow, NOAO, P.O. Box 26732, Tucson, AZ 85726, USA

R.A. WINDHORST

Department of Physics & Astronomy, Arizona State University, P.O. Box 871504, Tempe, AZ 85287-1504, USA

C. GRONWALL

Department of Astronomy, Pennsylvania State University, 525 Davey Laboratory, University Park, PA 16802

*Accepted by ApJ*

## ABSTRACT

We present a two-pronged approach to the formation of early-type galaxies, using a sample of 18 galaxies at  $0.5 \lesssim z \lesssim 1$  from the HST/ACS Ultra Deep Field and GRAPES surveys: 1) We combine slitless low resolution spectroscopy from the GRAPES dataset with simple models of galaxy formation to explore their star formation histories. 2) We also perform an analysis of their surface brightness distribution with the unprecedented details provided by the ACS superb angular resolution and photometric depth. Our spectroscopic analysis reveals that their stellar populations are rather homogeneous in age and metallicity and formed at redshifts  $z_F \sim 2 - 5$ . Evolving them passively, they become practically indistinguishable from ellipticals at  $z = 0$ . Also, their isophotal shapes appear very similar to those observed for nearby ellipticals, in that the percentages of disk and boxy galaxies at  $z \sim 1$  are close to the values measured at  $z = 0$ . Moreover, we find that the isophotal structure of  $z \sim 1$  early-type galaxies obeys the correlations already observed among nearby ellipticals, i.e. disk ellipticals have generally higher characteristic ellipticities, and boxy ellipticals have larger half-light radii and are brighter in the rest-frame B band. In this respect then, no significant structural differences are seen for ellipticals between  $z = 0$  and 1. Exception can be possibly made for the  $a_3/a$  parameter, which is larger at  $z \sim 1$  than usually measured at  $z = 0$ . The  $a_3/a$  parameter measures the deviations from a pure elliptical isophote which are not symmetric with respect to the galaxy center, as in the case of dust features and most notably of clumps. Blue clumps have been detected in nearly 50% of the  $z \sim 1$  early-type galaxies; their photometry is suggestive of young star clusters or dwarf irregulars if they are assumed to be at the same redshift as their host galaxies. We speculate that these clumps may represent recent accretion episodes, and that they could be a way to produce blue cores if their dynamical time is such for them to rapidly sink to the galaxy center.

*Subject headings:* galaxies: elliptical and lenticular, cD – galaxies: evolution – galaxies: formation – galaxies: stellar content

## 1. INTRODUCTION

Early-type galaxies form a remarkably homogeneous class of objects with a tight color-magnitude relation and a well defined Fundamental Plane. The small scatter in their colors (Bower, Lucey & Ellis 1992, Stanford, Eisen-

hardt & Dickinson 1998,) and mass-to-light (M/L) ratios (cf. Kelson et al. 2000) suggest that early-type galaxies formed the bulk of their stars at high redshift (i.e.  $z_F > 2 - 3$ , van Dokkum et al. 1998, Thomas et al. 2005). Since then, early-type galaxies continued to grow in mass at different paces: the more massive objects ( $> 10^{11} M_\odot$ ) appear to have increased their mass by less than 1% since  $z \simeq 1$ , while the less massive grew by 20 - 40% (cf. Chen et al. 2003, Bell et al. 2004, Cross et al. 2004, Conselice et al. 2005, Ferreras et al. 2005, Thomas et al. 2005, Treu et al. 2005). The detailed indices

Electronic address: pasquali@phys.ethz.ch  
Electronic address: ferreras@star.ucl.ac.uk  
Electronic address: panagia@stsci.edu  
Electronic address: edaddi@noao.edu  
Electronic address: Rogier.Windhorst@asu.edu  
Electronic address: caryl@astro.psu.edu

analysis of Thomas et al. (2005) shows that the more massive early-type galaxies are generally dominated by old stellar populations, enriched in the  $\alpha$  elements, while intermediate-age stars are found in early-type galaxies less massive than  $10^{10} M_{\odot}$  and with low  $[\alpha/\text{Fe}]$  ratios (cf. also Nelan et al. 2005). The combination of old stellar ages and high abundance ratios typical of massive early-type galaxies is suggestive of short formation timescales (i.e. 1 Gyr, Thomas, Greggio & Bender 1999), during which star formation occurred either with a flattened IMF (Thomas 1999), or through multiple bursts characterized by an enhanced star-formation activity. On the other hand, the spheroidal morphology and hot dynamics observed in these systems require major mergers, which result in a more extended dynamical history. The discrepancy in the star-formation and dynamical timescales may be reconciled through the *progenitor bias* (van Dokkum & Franx 2001), whereby about 50% of the present-day early-type galaxies were transformed from star-forming galaxies at  $z < 1$ , with the latter possibly having a constant star formation rate prior to their morphological transformation. The question, at this point, is whether the scaling relations derived for early-type galaxies at  $z < 1$  apply also to early-type galaxies at intermediate redshift. The high angular resolution of the Advanced Camera for Surveys (ACS) onboard HST allows for the very first time to measure the structural parameters of early-type galaxies in the UDF/GRAPES surveys with redshifts between 0.5 and 1.3. In this paper, we will discuss their star formation histories as derived from their GRAPES (GRISM ACS Program for Extragalactic Science, Pirzkal et al. 2004) spectra and their possible assembling histories as deduced from their isophotal structure in the UDF images (a concordance cosmology,  $\Lambda\text{CDM}$ ,  $\Omega_m = 0.3$ ,  $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ , is assumed hereafter).

The traditional approach to the formation of early-type galaxies is to study their stellar populations via spectrophotometric observables (see e.g. Worthey 1994; Trager et al. 2000; Bernardi et al. 2003), derive the age and metallicity distribution of their (unresolved) stellar content and finally reconstruct their star-formation history. This is the ultimate ingredient in distinguishing between different mechanisms for the formation of ellipticals, such as monolithic collapse (Eggen, Lynden-Bell & Sandage 1962) and hierarchical merging (Kauffmann et al. 1993). At the same time, the study of the morphology of early-type galaxies allows us to constrain their assembly history.

The advent of wide-field imaging surveys performed from the ground and with HST (e.g. Williams et al. 1996, Giavalisco et al. 2004, Beckwith et al. 2005) has proved that galaxy structure evolves with time, so that more distant galaxies are more peculiar than those in the local Universe on which the Hubble classification was originally based (Driver et al. 1995, Glazebrook et al. 1995, Abraham et al. 1996, Driver et al. 1998, Brichmann & Ellis 2000, Conselice et al. 2005). Furthermore, the morphology-density relation (Dressler 1980, Dressler et al. 1997) confirms that galaxy morphology depends strongly on environment, with early-type galaxies preferentially living in high density regions. Notice, here, that the morphological K-correction can affect the morphological classification, depending on the fading of the surface

brightness with wavelength. Windhorst et al. (2002, cf. also Colley et al. 1996) showed that early-type galaxies endure a significant decrease in their surface brightness from red to mid-UV wavelengths which could lead to a different classification and makes them almost undetectable at intermediate to high redshifts. Mid-type and star-forming galaxies appear in the mid-UV with a somewhat later type, while the majority of late-type and merging systems have a morphology little dependent on wavelength.

Galaxy shapes can be quantified into several “morphological parameters” (e.g. surface brightness profile, bulge-to-disk ratio, ellipticity, light concentration, asymmetry and clumpiness) which turn out to correlate with the galaxy star-formation rate, stellar mass, central black hole mass and merging history. For example, the light concentration of a galaxy varies with its luminosity, stellar mass, size and the mass of the central black hole (Caon et al. 1993, Graham et al. 1996, Bershadsky et al. 2000, Graham et al. 2001, Conselice 2003). Therefore, it depends on the past formation history of a galaxy. Asymmetry and clumpiness “measure” a more recent epoch in the evolution of a galaxy, since asymmetry is due to the presence of a merger and/or to tidal interactions, and clumpiness correlates with the degree of on-going star formation (Conselice 2003). These correlations allow us to decipher galaxy morphologies into mechanisms of galaxy formation and into time evolution of galaxy assembly.

Extensive imaging from the ground and with HST has revealed that a large fraction of nearby elliptical galaxies (Es) differ from a pure ellipse shape and do show substructures. For example, the ellipticity is not always constant with distance from the galaxy center. Di Tullio (1978, 1979) found that isolated Es are, on average, characterized by a decreasing ellipticity, while the mean ellipticity profile is either increasing or peaking with radius in Es located in galaxy clusters and groups. A change in the position angle of the galaxy major axis is often observed and explained as a projection effect of a triaxial structure whose axes ratios ( $b/a$  and  $c/a$ ) vary with radius (Galletta 1980, Franx 1988). About 60% of Es have “non-elliptical” isophotes, equally separated into *disky* or *boxy* (Bender et al. 1988, 1989). Disky Es are generally oblate rotators, while boxy Es span a variety of kinematical properties and are systematically bigger and brighter, suggesting that they formed via mergers (Naab, Burkert & Hernquist 1999). Almost half of the nearby Es, especially those in galaxy groups, show dust lanes which are aligned with the major or minor axis and thus believed to be byproducts of mergers (Lauer 1985, Sadler & Gerhard 1985, Ebneter & Balick 1985, van Dokkum & Franx 1995, Tran et al. 2001, Lauer et al. 2005, Martel et al. 2004). Elliptical galaxies surrounded by shells have also been observed and are thought to be the result of a collision with a disk galaxy (Malin & Carter 1983, Quinn 1984, Michard & Prugniel 2004). Finally, a blue core, possibly experiencing star formation, has been detected in  $\sim 30\%$  of nearby Es (Abraham et al. 1999, Papovich et al. 2003, Menanteau et al. 2001, Goto 2005) and nuclear stellar disks in  $\sim 60\%$  of early-type galaxies (van den Bosch et al. 1994, Rest et al. 2001, Lauer et al. 2005).

Our knowledge of the detailed structure of early-type

galaxies is mostly based on what has been observed so far in the local Universe. The purpose of this paper is to extend this study to early-type galaxies at intermediate redshift, so to describe in detail their structure - redshift correlation and to complement this with an accurate study of their stellar populations.

## 2. OBSERVATIONS

The Hubble Ultra Deep Field (UDF) is a survey of a  $3'.4 \times 3'.4$  field located in the Chandra Deep Field South, which was carried out with the Advanced Camera for Surveys (ACS) in four filters: F435W, F606W, F775W and F850LP (cf. Beckwith et al. 2005 for further details). We followed up these observations with ACS grism and slitless spectroscopy, as part of the GRAPES project which was awarded 40 HST orbits during Cycle 12 (ID 9793, PI S. Malhotra). The grism observations were taken at different epochs and at four different orientations in order to minimize the spectra contamination and overlapping from nearby sources. We have been then able to extract low resolution spectra ( $R \simeq 100$ , in the range 5500 Å to 10500 Å) for 5138 sources in the UDF field down to a limiting magnitude of  $m_{F850LP} \simeq 29.5$  in AB system. Readers are referred to Pirzkal et al. (2004) for a complete discussion of the data acquisition and reduction and for a description of the final GRAPES catalogue.

## 3. THE SAMPLE OF EARLY-TYPE GALAXIES

Our project aims at a combined analysis of the spectroscopic and morphological properties of early-type galaxies in the UDF. We thus selected from the GRAPES catalogue those sources with a high signal-to-noise ratio in their spectra, typically  $\text{SNR} \sim 10 - 20$  per pixel at  $\lambda = 8000$  Å; such values guarantee an accurate analysis of the observed spectral energy distributions (SEDs) in terms of stellar ages and metallicities. We found that this criterion imposes an apparent magnitude limit around  $m_{F775W} = 24.0$  in AB system. We then computed the concentration (C) and asymmetry (A) indices for the spectroscopically selected sample, in order to select those objects which lie in the parameter space defined by ellipticals (cf. Conselice 2003). Specifically, we derived the concentration and asymmetry parameters using the images taken in the F775W band, which roughly maps into a rest-frame *B* band at the redshifts reported in Table 1. We adopted for the concentration and asymmetry the same definitions and computing recipes given by Conselice (2003). We finally selected as *bona fide* early-type galaxies 18 objects, for which  $C \geq 3.2$  and  $A \leq 0.2$  in agreement with Conselice et al. (2005), and whose elliptical morphology was also visually confirmed. They are shown in Figure 1 together with the mean loci for nearby early-, late-type galaxies and mergers derived by Bershadsky et al. (2000) and Conselice et al. (2000). Table 1 shows the main properties of our final sample, including the UDF/GOODS ID numbers. Magnitudes are given in the AB system – taking the MAG\_BEST values from SExtractor (Bertin & Arnouts 1996). The optical magnitudes come from the UDF catalogue (Beckwith et al. 2005). Three of the galaxies in our sample (J033229.22-274707.6, J033240.33-274957.0, J033244.09-274541.5) are not in the UDF field of view although their

spectra were detected in the GRAPES survey. Their images were extracted from the GOODS data (Gialavalisco et al. 2004). UDF4527 and J033229.22-274707.6 have been also detected by Chandra in the soft X-rays, with a flux of  $3 \times 10^{-17}$  erg s $^{-1}$ cm $^{-2}$  and  $8 \times 10^{-17}$  erg s $^{-1}$ cm $^{-2}$  respectively (Alexander et al. 2003, Giacconi et al. 2002).

The last two columns of Table 1 give the spectroscopic redshifts derived with the ACS grism and with the VLT (FOR2: Vanzella et al. 2005, VIMOS: Le Fèvre et al. 2004). The redshifts obtained from the ACS grism have been measured by matching templates corresponding to a range of stellar populations to the low resolution GRAPES spectra. For those galaxies with an accurate VLT redshift, our grism estimates agreed to  $\Delta z / (1 + z) \lesssim 0.01$ , significantly better than any photometric redshift estimate (see e.g. Mobasher et al. 2004). The *bona fide* early-type galaxies span the redshift range between  $z \simeq 0.5$  and 1.3, with about 80% of the sample being distributed between  $z \simeq 0.5$  and  $z \simeq 0.8$ . Notice here that 39% of the *bona fide* early-type galaxies are clustered around  $z \simeq 0.665$ . This is a well known spike, since it has been detected in the K20 sample of ellipticals (Cimatti et al. 2002), in the Chandra Deep Field South (Gilli et al. 2003) and in the VLT spectroscopic follow-up of GOODS (Vanzella et al. 2005). These surveys indicate that the galaxies at  $z \simeq 0.67$  constitute a loose structure rather than a cluster.

## 4. STAR FORMATION HISTORIES

Our first approach in the study of the sample involves the star formation histories. In order to break the age-metallicity degeneracy, we combined models of galactic chemical enrichment and the low resolution slitless spectroscopy from the GRAPES data. We took extra care in avoiding contamination from nearby sources, which – in some cases – required the use of data corresponding to a single orientation.

In order to make a robust assessment of the ages and metallicities of the unresolved stellar populations, we decided to use two different models to describe the star formation histories, as described below. Briefly, the models depend on a reduced set of parameters, which uniquely determine a star formation history (SFH), i.e. a distribution of stellar ages and metallicities. This SFH is used to combine simple stellar populations from the models of Bruzual & Charlot (2003) and obtain a spectral energy distribution (SED) assuming a Salpeter IMF. Each choice of SFH is compared with the data via a  $\chi^2$  test. We explore a wide volume of parameter space in order to infer robust constraints on the ages and metallicities of the stellar populations in these galaxies. This comparison requires a careful process of degrading the synthetic SED (resolution  $R \sim 2000$ ) to the (variable) resolution of the GRAPES spectra. Special care must be taken with respect to the change of the PSF with wavelength, which results in both an effective degradation of the spectral resolution as a function of wavelength, and a different net spectral resolution with respect to the size of the galaxy. The latter requires the spectral resolution to be treated as a free parameter, to be chosen by a maximum likelihood method. We found that this parameter is not degenerate with respect to those describing the star formation history, and mostly results in a global shift of the likelihood. The two models chosen to describe the

build-up of the stellar component are as follows:

**Model #1 (EXP):** We take a simple exponentially decaying star formation rate so that each history is uniquely parameterized by a formation epoch – which can be described by a formation redshift ( $z_F$ ), a star formation timescale ( $\tau_*$ ) and a metallicity ( $[m/H]$ ) which is kept fixed at all times.

**Model #2 (CSP):** We follow a consistent chemical enrichment code as described in Ferreras & Silk (2000). The model allows for gas infall and outflows and keeps the star formation efficiency as a free parameter. The metallicity evolves according to these parameters, using the stellar yields from Thielemann, Nomoto & Hashimoto (1996) for massive stars ( $> 10M_\odot$ ) and van den Hoek & Groenewegen (1997) for intermediate mass stars. The free parameters are the star formation efficiency, the fraction of gas ejected in outflows, the formation epoch, and the timescale for the infall of gas.

Figure 2 shows the best fits and the observed SEDs<sup>1</sup>. The error bars represent the observations and the solid line correspond to the best fits for the CSP model. Figure 3 shows the ages and metallicities of the best SFHs for each galaxy. Average and RMS scatter for age and metallicity are shown as dots and error bars, respectively. Notice the metallicity in the EXP models do not have an error bar as this model assumes a fixed value of metallicity for each SFH. The lines in the bottom panel give the age of the Universe as a function of redshift (solid line) and the ages at that redshift corresponding to a formation epoch of (dashed, from top to bottom)  $z_F = \{5, 3, 2\}$ . Analogously to the analysis of the surface brightness distribution shown in the next sections, we find that the stellar populations in these galaxies are dominated by old, passively evolving stars, formed at redshifts  $z_F \gtrsim 2$ . Notice that the EXP model tends to underestimate the stellar ages. This model assumes a constant metallicity, which often forces older populations to have higher metallicities, which is compensated by the age-metallicity degeneracy by shifting the average towards younger ages. The more consistent model (CSP) generates a more realistic distribution of metallicities and we should expect it to be closer to the true populations. Nevertheless, the difference between these models is rather small, and we can safely reject a significant star formation at  $z \sim 1 - 1.5$  in *all* galaxies. These models also give galaxy mass estimates in the range  $9.5 \leq \log(M/M_\odot) \leq 11.5$ .

The star formation histories that give the best fit can be used to predict the evolution of the color-magnitude relation (CMR). Figure 4 shows the rest-frame  $U - V$  vs.  $M_V$  CMR of our sample. The filled dots are simple translations of the observed photometry to restframe  $U - V$  (i.e. a K correction) whereas the open dots evolve these galaxies to zero redshift, in order to compare our galaxies with local samples. Two characteristic error bars are shown. We plot Coma cluster galaxies from Bower, Lucey & Ellis (1992) and the dashed lines rep-

resent the linear fit and scatter. The Figure shows that our sample of early-type galaxies is fairly consistent with the stellar populations found in local elliptical galaxies. Notice that the passive fading of the brightest galaxies puts them at  $z = 0$  within the luminosity range of the brightest local ellipticals. The available information also allows us to explore a projection of the Fundamental Plane, namely the Kormendy relation (Kormendy 1977). Figure 5 shows this relation (filled dots with error bars) compared to Coma cluster galaxies (squares; Jørgensen, Franx & Kjaergaard 1995), a sample of distant early-type galaxies in the HDF (grey dots; Fasano et al. 1998), the LBDS radio elliptical galaxies at  $z \sim 1.5$  (open stars; Waddington et al. 2002) and the early-type galaxies at  $z \sim 2$  studied by Daddi et al. (2005, open dots). The translation from the observed photometry to  $\mu_B$  involves a K correction as well as the  $(1+z)^4$  cosmological dimming. The offset  $\Delta\langle\mu_B\rangle \sim 1.5$  mag between our sample and Coma (as well as between Coma and the HDF sample and the LBDS radio galaxies) is compatible with the expected fading of the stellar populations. Notice that both the GRAPES and the HDF sample lack  $r_{hl} \sim 10$  kpc galaxies, most likely due to the limited volume sample in the HDF and UDF surveys (the  $r_{hl}$  of the *bona fide* early-type galaxies was measured in the F775W filter, corresponding to the restframe B band at  $< z > \simeq 0.7$ ). No *bona fide* early-type galaxy at  $z \simeq 1$  is as compact as those studied by Daddi et al. (2005) at  $z \sim 2$ .

## 5. ISOPHOTAL SHAPES

We used IRAF ELLIPSE routine (Jedrzejewski 1987) to measure the isophotal parameters of the *bona fide* early-type galaxies in the F606W, F775W and F850LP bands. At a mean redshift of 0.7 (cf. Table 1), these filters sample the restframe U, B and V bands. ELLIPSE allows the isophote center, ellipticity and position angle to vary at each iteration. Once a satisfactory fit is found, it computes the sin and cos  $3\theta$  and  $4\theta$  terms, which describe the deviations of the isophote from pure ellipse by means of the following Fourier expansion:

$$\Delta r(\theta)/r(\theta) = (a_3/a)\cos(3\theta) + (a_4/a)\cos(4\theta) + (b_3/b)\sin(3\theta) + (b_4/b)\sin(4\theta)$$

where  $\theta$  is the position angle,  $r$  the distance from the galaxy center and  $a$  and  $b$  the semi-major and semi-minor axes. The  $a_4/a$  and  $a_3/a$  parameters, where  $a$  is the semi-major axis of the isophote, measure the deviations of the isophote from a pure ellipse. In particular,  $a_4/a$  measures the deviations symmetrical with respect to the galaxy center (i.e. found along the isophote every  $90^\circ$ ), while  $a_3/a$  the non-symmetrical deviations occurring at every  $120^\circ$ .

We ran ELLIPSE leaving free the position of the center, the ellipticity (defined as  $1 - b/a$ , where  $a$  and  $b$  are the projected major and minor axes of a galaxy) and the position angle of the major axis and assuming a logarithmic step along the semi-major axis. The results are shown for each individual galaxy in Figures 11a - 11r (of which Figures 11b - 11r are available in the electronic edition of the Journal), where the surface brightness, ellipticity, position angle (PA) of the major axis, the  $a_3/a$  and  $a_4/a$  parameters are plotted for each filter as a function of distance from the galaxy center. These radial profiles extend to about  $1''.2$  from the galaxy center in order to avoid overlap with nearby objects.

<sup>1</sup> The spectra of J033229.22-274707.6 and UDF6027 could resemble those of dusty ULIRGS (cf. Moustakas et al. 2004). The CAS morphology of these two galaxies is still consistent with them being ellipticals although somewhat close to the transition edge between early-type galaxies and ULIRGs. The low resolution of the ACS grism certainly helps in smoothing the spectra and decreasing the Balmer break.

The ellipticity of the *bona fide* early-type galaxies show a variety of radial dependencies: increasing outward (i.e. J033229.22-274707.6, UDF9264, UDF2322, UDF8, UDF153, J033240.33-274957.0, J033244.09-274541.5), decreasing outward (UDF3677), peaked (i.e. UDF8316, UDF6747, UDF68, UDF4587, UDF4527, UDF2107, UDF5177), nearly flat (i.e. UDF2387, UDF1960, UDF6027). In the case of UDF1960, UDF4587 and UDF5177, the radial profile of the ellipticity is strongly perturbed by more compact (and usually bluer) sources projected onto the main body of these galaxies. These features are somewhat dependent on wavelength, since the ellipticity profiles appear to be smoother and to extend to lower values in the restframe V band (i.e. the F850LP filter). Di Tullio (1978, 1979) analysed a sample of 75 nearby ellipticals and found that the radial variation of a galaxy ellipticity is different for different environments. Indeed, ellipticals with increasing ellipticity are found in galaxy clusters and groups, and systems with peaked ellipticity also in galaxy pairs. Ellipticals with constant or decreasing ellipticity are preferentially isolated, although are also observed in galaxy clusters, groups and pairs. Di Tullio (1978, 1979) suggested that an increasing ellipticity with radius could be a signature of merger and/or tidal interactions with a companion galaxy.

The position angle of the *bona fide* early-type galaxies is seen to generally vary with radius. The galaxies UDF1960, UDF3677, UDF68 and UDF8 show significant troughs in their PA profiles, which are likely due to the presence of dust features and compact sources projected onto the main body of these galaxies (cf. Sect. 7). Galletta (1980) explained the radial PA variations as the effect of the projection of a triaxial structure with the axes ratios ( $b/a$  and  $c/a$ ) varying with radius, onto the plane of the sky. Di Tullio (1979, 1980) proposed that PA variations larger than  $10^\circ$  could be relics of past mergers. Isophotes can locally vary from a simple ellipse shape: if this variation occurs every  $90^\circ$  along the isophote and from the galaxy center, it indicates that the isophote has either a disk or a boxy shape. This variation is quantitatively measured with the  $a_4/a$  parameter; disk isophotes have  $a_4/a > 0$ , boxy isophotes are characterized by negative values of  $a_4/a$ , while elliptical isophotes have  $a_4/a = 0$  (Lauer 1985, Bender et al. 1988, 1989). In our sample of *bona fide* early-type galaxies, the radial variation of the  $a_4/a$  parameter is quite dependent on wavelength, except for the galaxies UDF1960, UDF3677, UDF8316, UDF6747, UDF68, UDF2322, UDF6027 and UDF4527. The early-type galaxies which are clearly disk ( $a_4/a \gg 0$ ) at any wavelength are: UDF2387, UDF6747, UDF4527, UDF2107 and UDF5177, while the only clear boxy galaxy is UDF3677. More elliptical galaxies (with no significant variations at any wavelength) are J033229.22-274707.6, UDF1960, UDF8316, UDF2322, UDF6027, J033240.33-274957.0. The galaxies mentioned above may show troughs or sharp peaks in the radial profile of their  $a_4/a$  parameter, which can be produced by dust features and compact sources projected onto the main body of these galaxies. UDF153 is the only galaxy that becomes boxy only in the restframe B and V bands (i.e. F775W and F850LP filters). Other galaxies feature a radial change from disk to boxy (and vice-versa), such as UDF9264, UDF68, UDF8, UDF4587

and J033244.09-274541.5. In particular, UDF8 changes from elliptical to disk in the restframe U and B bands (i.e. F606W and F775W filters), while it changes from boxy to disk in the restframe V band (F850LP filter). The radius at which this change occurs is in the area where the PSF still dominates the light distribution of the galaxy (cf. Sect. 5).

A second parameter,  $a_3/a$ , measures the deviation of an isophote from a perfect ellipse, every  $120^\circ$  along the isophote and from the galaxy center. This variation is mostly induced by the presence of dust features and clumps in the galaxy. Among the *bona fide* early-type galaxies, we detect  $a_3/a \neq 0$  in UDF2387, UDF1960, UDF3677, UDF8316, UDF6747, UDF8, UDF6027, UDF153, UDF4587, UDF4527, J033244.09-274541.5, UDF2107 and UDF5177.

The measured isophotal parameters exhibit radial gradients that do not appear to significantly depend on the wavelength at which the galaxies were imaged. Given the caveat on the effects of the morphological K-correction (Windhorst et al. 2002), this confirms the robustness of the criteria we used to select our sample of *bona fide* early-type galaxies.

Bender et al. (1988, 1989) analysed a sample of nearby ellipticals and concluded that they equally split among boxy systems (30%), disk galaxies (30%) and ellipticals with no significant deviations (30%). In particular, disk ellipticals have ellipticity larger than 0.25 and behave as oblate rotators, while boxy systems have ellipticities in the range between 0.15 and 0.45 and display a large variety of kinematical properties. In addition, boxy ellipticals are on average bigger with half-light radii larger than 5 kpc and brighter than disk systems in the B band, at radio wavelengths and in the X-rays. Starting from the idea whereby ellipticals can originate from the merger of two disk galaxies (Toomre & Toomre 1972), N-body simulations of disks collisions predict kinematic and photometric properties of the merger remnants very similar to those observed for elliptical galaxies, i.e. decoupled cores and isophotal shapes (cf. Barnes 1992, Hernquist 1992, Naab, Burkert & Hernquist 1999, Naab & Burkert 2003). Khochfar & Burkert (2005) have shown that the morphology of the merging galaxies and any subsequent episode of gas infall are critical in determining the shape of the merger remnant. Boxy ellipticals can be reproduced via a merger of two equally-massive disk galaxies or a major merger of two early-type galaxies. Disk ellipticals are mainly formed through an unequal-mass merger of two disk galaxies or late gas infall. In the hierarchical framework equal-mass mergers are expected to be much rarer than the unequal-mass, although the observational selection seems to go in the opposite direction (i.e. the ratio of boxy to disk galaxies at  $z \simeq 0$  is nearly one).

## 6. AVERAGE ISOPHOTAL PARAMETERS

Given that the *bona fide* early-type galaxies in the UDF/GRAPES survey are at higher redshift than the nearby ellipticals studied by Bender et al. (1988, 1989), it is interesting to check whether the isophotal parameters show any change with redshift. We therefore computed for each galaxy the characteristic ellipticity and isophotal twist  $\Delta PA$ , the mean  $a_3/a$  and  $a_4/a$  parameters ( $\langle a_3/a \rangle$  and  $\langle a_4/a \rangle$ ) following the prescriptions of Bender et al.

(1988) and in the F850LP filter which for the observed redshifts corresponds to the restframe V and R bands used by Bender et al. Briefly, we averaged the  $a_3/a$  and  $a_4/a$  values measured at  $R_i \leq r \leq R_o$  weighted by their errors and by the counts in the isophote at each isophotal mean radius  $r$ . Here,  $R_o$  is 1.5 times the half-light radius  $r_{hl}$  derived from the  $\mu_{F850LP}$  radial profile and  $R_i$  is the radius corresponding to 3 times the FWHM of the PSF in the F850LP filter, chosen to avoid the central part of a galaxy where the light distribution is perturbed by the PSF. While  $R_i$  is 12 pixels, i.e.  $0''.36$ ,  $R_o$  varies up to  $1''.1$ . The isophotal twist was measured as a difference in PA between  $R_i$  and  $R_o$ , while the characteristic ellipticity was taken equal to the ellipticity at the half-light radius in case of increasing, decreasing or flat radial profiles, or equal to the peak ellipticity in case of peaked radial profiles. The final results are reported in Table 2, where the  $\langle a_3/a \rangle$  and  $\langle a_4/a \rangle$  values have been multiplied by 100. Since its  $R_o$  is comparable to  $R_i$ , we could not determine the isophotal parameters of J033240.33-274957.0 and UDF68.

As pointed out by Odewahn et al. (1997), the S/N ratio in the images can affect the measurement of the galaxy projected  $a$  and  $b$  axes and hence its ellipticity, in the sense that galaxies with a lower S/N value are measured to be rounder than real. Therefore, to test the reliability of the isophotal parameters derived for our sample, we measured the ellipticity,  $\langle a_3/a \rangle$  and  $\langle a_4/a \rangle$  parameters for a number of simulated ellipticals. The simulations were performed (cf. Appendix A) using a Sérsic profile (Sérsic 1968) with index between 3 and 4, and assuming the S/N ratio and the integrated magnitude of the observed data. The simulated images were then convolved with the ACS PSF in the F775W filter. We were able to recover for the simulated images not convolved by the PSF isophotal parameters very close to their input values, thus indicating that the S/N ratio of the data does not significantly affect our measurements. We also found that the ACS PSF underestimate the ellipticity by  $\simeq 10\%$ , overestimates  $\langle a_4/a \rangle$  by about 0.2 with respect to the null input value and introduces a scatter of about 0.2 in  $\langle a_3/a \rangle$  around the input  $a_3/a = 0$ . used for the simulations.

The isophotal parameters of the *bona fide* early-type galaxies (black filled circles) are plotted in Figure 6 as a function of the galaxy half-light radius  $r_{hl}$  (measured in the F850LP filter, corresponding to the restframe V band at  $< z > \simeq 0.7$ ), total absolute magnitude in restframe B band  $M_B$ , and compared with the sample of Bender et al. (1988, 1989) represented with grey, filled circles. It has to be specified that the  $M_B$  values were derived from the best-fitting, model spectra (cf. Sect. 4), also K-corrected. The magnitude range (hence stellar mass) of the nearby ellipticals is nicely sampled by the *bona fide* early-type galaxies, and this makes the comparison of the two galaxy samples in their isophotal parameters even more robust. We chose to correct the ellipticity and the  $\langle a_4/a \rangle$  parameter for the PSF, i.e we increased ellipticities by 10% and decreased  $\langle a_4/a \rangle$  by 0.15. Overall, the *bona fide* early-type galaxies confirm the trends traced by nearby ellipticals:

- i) boxy ellipticals tend to be bigger (as shown in Figure 6a).
- ii) Rounder ellipticals show smaller deviations from pure

ellipses, while larger deviations are seen in galaxies with ellipticity  $\geq 0.25$ . Disky ellipticals have  $0.15 \leq \text{ellipticity} \leq 0.6$ , while boxy ellipticals  $0.15 \leq \text{ellipticity} \leq 0.45$ . The deviant galaxies in our sample are UDF1960 and UDF6747 (cf. Figure 6b). The ellipse fitting of UDF1960 is affected by the presence of a blue clump (a probable lens candidate, cf. Sect. 7.4) at a distance of  $0''.4$  from the galaxy center, and by the disk-like substructure in UDF6747 (cf. Figure 10).

- iii) Boxy ellipticals tend to be brighter than disk systems (see Figure 6c).

Although not shown in Figure 6, the isophotal twist increases at lower ellipticity because of the uncertainties in determining the major axis PA of a rounder galaxy. We can thus conclude that ellipticity and  $\langle a_4/a \rangle$  do not significantly change with redshift up to  $z \simeq 1$ . About half of the *bona fide* early-type galaxies are characterized by values of  $\langle a_3/a \rangle$  larger than the range spanned by nearby ellipticals. Their large  $\langle a_3/a \rangle$  values are likely due to the presence of dust features (e.g. UDF2387 and UDF3677, cf. Sect. 7.1) or blue clumps close to the galaxy center (e.g. UDF4587 and UDF2107, cf. Sect. 7.3; UDF1960 with an overlapping lens candidate is a special case, cf. Sect. 7.4). When dust features, disk-like substructures and blue clumps are masked out, the isophote fitting gives significantly smaller values of the mean  $\langle a_3/a \rangle$  parameter, which then falls in the range spanned by nearby ellipticals.

## 7. SUBSTRUCTURES IN THE UDF EARLY-TYPE GALAXIES

We used the outputs of the ELLIPSE routine to build model images of the *bona fide* early-type galaxies, and subtracted them from the observed frames in order to look for residual substructures. In the cases where substructures were detected, we re-ran ELLIPSE masking out the substructures and adjusted the mask every run until the extinction map of each galaxy reached a minimum value of  $\simeq \pm 0.05$  mag in the unmasked pixels. The extinction map was computed from the ratio of the observed image to the ELLIPSE model image with the formula:  $2.5 \times \text{Log}(\text{Observed}/\text{Model})$ . We applied this procedure to the images obtained in the F606W and F775W filters, which are characterized by a better S/N ratio and PSF with respect to the F435W and F850LP filters.

In this way, we could detect dust features, disks and clumps in a significant fraction of our sample. Six galaxies do not show either a disk substructure nor dust features: J033229.22-274707.6, J033240.33-274957.0, J033244.09-274541.5, UDF2322, UDF6027 and UDF8316. J033244.09-274541.5 is shown in Figure 9: a close-up of the central region ( $1''.2 \times 1''.2$ ) in the F775W filter is in the left-hand panel in units of counts/s, and the extinction map of the same region in units of magnitude (mean residual value  $\simeq \pm 0.05$  mag/pix) in the right-hand side panel. The (V-I) color maps of the six featureless galaxies together with the remainder of our sample are shown in Figures 7 and 8. The (V-I) color maps were obtained using Voronoi tessellations, so that pixels were binned in order to reach a chosen and constant S/N ratio per bin (cf. Cappellari & Copin, 2003). In our case, the targeted S/N ratio corresponds to 0.05 mag the color uncertainty.

### 7.1. Dust features

Dust features have been resolved in UDF2387 and UDF3677 (see Figure 9). They appear as white stripes in the extinction maps (right-hand side panels of both Figures). Here, grey indicates a residual of about 0.05 mag and white an extinction  $A_{F775W}$  ( $\simeq$  restframe  $A_B$ ) up to  $\simeq 0.25$  mag in UDF2385 and 0.18 mag in UDF3677. These values are very similar to what found for dusty nearby ellipticals at  $z \simeq 0$  by Sadler & Gerhard (1985).

### 7.2. Disks

UDF4527 and UDF6747 are characterized by a disk-like structure that is aligned with the galaxy major axis and has a full extension of about  $1''$  (corresponding nearly to the half-light radius of each galaxy). These features are shown in Figure 10, where they turn out to be in emission (black pixels) by  $\sim 0.10$  mag/pix in UDF4527 and  $\sim 0.25$  mag/pix in UDF6747. In these images, light grey levels indicate residuals of about  $\pm 0.05$  mag. In the (V-I) [restframe (B-V)] color map of UDF4527 and UDF6747, the disk turns out to be as red as the galaxy and the two should thus be coeval. Such a similarity is known to be common among S0 galaxies, because of either small age differences (Peletier & Balcells 1996) or a larger amount of dust in the disk (Michard & Poulain 2000).

### 7.3. Blue clumps

As seen in Figures 7 and 8, a large fraction of *bona fide* early-type galaxies shows compact, blue clumps at different distances (up to  $2''$ ) from the galaxy center. UDF6747, in particular, is surrounded by a cluster of blue clumps at one edge of its major axis (cf. Figure 8). Figure 10 highlights this feature in UDF153. We could measure the integrated magnitude of these clumps in the four filters of the UDF after having subtracted the ELLIPSE model image from the observed frame of each galaxy. It turns out that these clumps are very faint, with magnitudes between 30.9 and 26.1, and contribute to the total flux of the host galaxy at a  $\leq 4\%$  level. Hence, these clumps do not contribute significantly to the grism spectra of the *bona fide* early-type galaxies. What is the nature of the blue clumps? The simplest explanation would be that they are sources at high redshift, as in the case of UDF1960, where the blue clump (cf. Figure 7) is a lens candidate (Fassnacht et al. 2004). On the other hand, given the number of objects detected in the UDF field and the size of the galaxies in our sample, we have estimated a probability of about 30% for objects to overlap. A more intriguing possibility is that the most compact of these clumps are physically associated with the *bona fide* early-type galaxies, and, as such, may be HII regions or young star clusters or satellite dwarf galaxies. To test this hypothesis, we ran the Hyperz code (Bolzonella et al. 2000) on the set of magnitudes measured for each clump, assuming that the clump is at the same redshift of the galaxy in which it has been observed. The reduced  $\chi^2$  of the fits indicates that some of these clumps could be consistent with the above assumption, and they could be young (few  $10^6$  - few  $10^8$  yrs) systems in the range  $-12 \leq M_V \leq -16$ . Though speculative, these results are in good agreement with the study of O'Dea et al. (2001) who found four star-forming regions in the double-double

radio galaxy 3C 236, and of Yang et al. (2004) who detected blue compact sources in five E+A galaxies at  $z \simeq 0.1$ .

### 7.4. Peculiar features

UDF68, UDF1960, UDF2107 and UDF5177 display in their extinction map some arc-like features in emission. An example is given in Figure 10 for UDF5177. With the data in hand, it is not possible to establish whether these features are real substructures or, for example, lensed sources at high redshifts overlapping with the UDF early-type galaxies. Fassnacht et al. (2004) have identified only one lens candidate among the *bona fide* early-type galaxies, which lies nearby UDF1960. We suggest that UDF68 has also lensed sources projected onto its main body, given that the morphology of its (V-I) map is quite similar to UDF1960. The arc feature in UDF5177 is apparently offset from the center of the light distribution of the galaxy and could be interpreted as the remnant of an accreted dwarf galaxy which went tidally disrupted.

### 7.5. Blue cores

Two galaxies in our sample exhibit a blue core: UDF5177 and UDF6027 (cf. Figure 8). About 30% of nearby ellipticals have a blue core, which is usually believed to have experienced recent star formation (Abraham et al. 1999, Papovich et al. 2003, Menanteau et al. 2001, Goto 2005). Menanteau et al. (2004) found a blue-core elliptical at  $z = 0.624$  which seems to host an AGN. In our sample, UDF5177 and UDF6027 are at  $z = 0.50$  and  $1.32$ , respectively.

## 8. CONCLUSIONS

According to the current paradigm of galaxy formation, ellipticals form through the merger of galaxies. As shown by Kauffmann & Charlot (1998) this predicts a fairly extended assembly history, which is difficult to reconcile with the old stellar age and the  $\alpha$ -enhanced chemistry of these galaxies. Because of their merger history, one would expect the isophotal morphology of early-type galaxies to be more distorted at high redshift compared to nearby ellipticals. In this respect, the UDF survey is an unique dataset which allows, for the very first time, to measure accurate isophotal parameters for 18 *bona fide* early-type galaxies between  $z \simeq 0.5$  and  $1.3$ . At the same time, the GRAPES survey provides for these same objects high S/N grism spectra, which are used to determine the their stellar age and formation redshifts. The population of  $z \sim 1$  early-type galaxies in the UDF is clearly old. The analysis of their spectra indicates that their stellar populations are rather homogenous in age and metallicity and formed at redshifts  $z_F \sim 2 - 5$ . Evolving them passively, they become indistinguishable from the ellipticals observed in the Coma cluster [cf. also Daddi et al. (2005) for a sample of UDF early-type galaxies at  $z \sim 2$ ]. As discussed by Hamilton (1985), spectroscopically-selected early-type galaxies are generally seen long after (i.e.  $\gg 1$  Gyr) the last major merger. After such a merger, these objects quickly settle back to their quasi steady-state, and will then look for a much longer time like non-evolving ellipticals: it is during this phase that we select and observe them now (cf. also van Dokkum & Franx 2001).

The  $z \sim 1$  early-type galaxies in the UDF are morphologically very similar to nearby ellipticals. They separate into disk galaxies ( $\sim 44\%$  with  $\langle a_4/a \rangle > 0.5$ ), boxy galaxies ( $\sim 31\%$  with  $\langle a_4/a \rangle < -0.5$ ) and systems with no significant deviation from a pure ellipse ( $\sim 25\%$  with  $-0.5 \leq \langle a_4/a \rangle \leq 0.5$ ). Given the small field of view of the UDF and therefore the small sample, these fractions can be taken as in good agreement with the results of Bender et al. (1988, 1989) who found  $\sim 30\%$  of the nearby ellipticals to be disk,  $\sim 30\%$  boxy and  $\sim 30\%$  with no significant deviations. The slight overabundance of disk *vs* boxy ellipticals in our sample could be explained by their environment. Indeed, VLT spectroscopic follow-up has detected three large-scale structures in the UDF (at  $z \simeq 0.67, 0.73, 1.10$  and  $1.61$ , Vanzella et al. 2005, see also Cimatti et al. 2002 and Gilli et al. 2003) which make the UDF an over-dense region at any of these three redshifts. In over-dense regions, the N-body simulations of Khochfar & Burkert (2005) predict a higher number of disk *vs* boxy ellipticals. The isophotal structure of the *bona fide* early-type galaxies obeys the correlations already observed among nearby ellipticals: *i*) disk ellipticals have generally higher characteristic ellipticities; *ii*) boxy ellipticals have larger half-light radii and are brighter in the B band. In this respect, the isophotal shape of ellipticals does not significantly change between  $z = 0$  and  $1$ . This suggests that these galaxies either do not merge significantly in their evolution towards  $z = 0$ , or mergers at later redshifts do not vary the percentages of disk and boxy ellipticals. In this respect, Schade et al. (1999) suggested that merging is not required since  $z = 1$  to produce the present-day space density of ellipticals. The  $\langle a_3/a \rangle$  parameter appears to be larger than measured by Bender et al. for nearly one third of the *bona fide* early-type galaxies. This deviation is likely due to the presence of dust features (in  $\sim 10\%$  of the sample, a percentage a factor of 4 smaller than observed at  $z = 0$ ) and most notably of clumps close to the galaxy center. Blue (in the (V-I) color) clumps have been detected in about 50% of the *bona fide* early-type galaxies, at different distances from the center (up to  $2''$ , in the galaxy outskirts). From their photometry and under the assumption that they are at the same redshift as their host galaxies, we have derived ages (from few  $10^6$  to few  $10^8$  yrs) and luminosities ( $-12 \leq M_V \leq -16$ ) consistent with them being either young star clusters or dwarf irregular galaxies. They appear very similar to the blue compact sources observed in five E+A galaxies at  $z \simeq 0.1$  by Yang et al. (2004), who suggested these sources to be young star clusters. On the other hand, the hypothesis of being dwarf galaxies would put these blue clumps in the right mass range to explain the small mass increase undergone by ellipticals more massive than  $10^{11.5} M_\odot$  (i.e. these galaxies seem to have grown less than 1% of their mass since  $z \simeq 1$  according to the results of Treu et al. 2005, or less than 5% as discussed by Schade et al. 1999). Gas accretion is also invoked to explain the presence of four star-forming regions in the double-double radio galaxy 3C 236; their age difference would compare with the timescale of the nuclear activity in this galaxy and hence would set time constraints on the infall of gas experienced by 3C 236 (O’Dea et al. 2001).

To add more speculation to the picture, since the blue clumps in our sample galaxies are distributed at differ-

ent distances from the galaxy center, we could imagine that they have been accreted and may possibly sink to the galaxy center, forming a blue core as observed in UDF5177 at  $z \simeq 0.5$ . We might then be facing the build-up of early-type galaxies at  $z \simeq 1$ , similarly to what observed by Chen et al. (2003), Bell et al. (2004), Cross et al. (2004), Conselice et al. (2005) and Ferreras et al. (2005). Clearly, a dedicated spectroscopic campaign would be needed to unveil the true nature of these blue clumps.

In the sample of *bona fide* early-type galaxies, UDF153 is particularly remarkable, since it is surrounded by a faint and incomplete shell. As shown by Quinn (1984), such a shell is the relic of a relatively recent merger with a smaller *disk galaxy* and can be observed for about 1 Gyr. The high phase-space density of a disk galaxy is indeed required to produce such a spatially confined substructure as the observed shells are. The GRAPES spectrum of UDF153 indicates a redshift of 0.98, suggesting that the merger possibly occurred at  $z \geq 1$ . This would also imply that some disk galaxies were already in place at  $z \geq 1$ , despite observations which show that the fraction of regular disk galaxies decreases at  $z > 1$  (cf. Abraham et al. 1996, Conselice et al. 2003).

N-body simulations (cf. Khochfar & Burkert 2005, Jesseit et al. 2005) indicate that the boxyness/diskyness shape of an elliptical galaxy is a long-lived signature of a merger if no further disturbances (i.e. late gas infall and/or subsequent mergers) occur. Indeed, the stellar orbits which forge the  $a_4/a$  parameter are stable and in equilibrium. Therefore, if the system stays unperturbed after the major merger which formed it, the “lifetime” of its isophotal shape is comparable with the age of its stellar population. Moreover, the isophotal shape of the merger remnant depends on the mass ratio of the merger and on the morphology (i.e. stellar orbits) of the merging galaxies. Given that the fractions of disk and boxy systems among the *bona fide* early-type galaxies is similar to the one observed at  $z = 0$ , and modulo the small size of their sample, the *bona fide* early-type galaxies can be used to trace the mass ratio of mergers as a function of redshift and the lifetime of the different isophotal shapes taken by ellipticals in N-body simulations like those of Khochfar & Burkert.

A last comment is devoted to the masses of the *bona fide* early-type galaxies. Our spectroscopic analysis indicates masses between  $10^{9.5}$  and  $10^{11.5} M_\odot$  in stars, which likely formed at about  $z \simeq 3$ . At this redshift, such masses are consistent with Lyman-break galaxies (LBG, Steidel et al. 1996, Shapley et al. 2001, Papovich et al. 2001), SCUBA galaxies (Bertoldi et al. 2000, Smail et al. 2002, Genzel et al. 2003, Webb et al. 2003, Chapman et al. 2003, 2004) and bright radio galaxies (van Breugel et al. 1998, Dey et al. 1997). These galaxies are all observed undergoing intense star formation with a rate varying from  $\sim 90 M_\odot \text{yr}^{-1}$  (in LBGs, Shapley et al. 2001) up to  $500 M_\odot \text{yr}^{-1}$  (in SCUBA sources, Genzel et al. 2003) and  $1000 M_\odot \text{yr}^{-1}$  (in high-redshift radio galaxies, Dey et al. 1997). Such a star-formation rate is able to enrich the galactic ISM up to a metallicity of about  $0.4 Z_\odot$  (Pettini et al. 2001), a value very similar to what found in this paper for the *bona fide* early-type galaxies. Such starbursts are also likely to produce the  $\alpha$ -element enhancement typical of elliptical galaxies (Trager et al.



2000). These similarities in mass and metallicity add further evidence to the evolution of vigorously star-forming galaxies at high redshift into  $z \simeq 1$  ellipticals.

We would like to thank an anonymous referee for useful comments that improved the paper. We also thank A. Burkert, F.C. van den Bosch and C. Porciani for valuable discussions. ED acknowledges support from NASA through the Spitzer Fellowship Program, under award

1268429. The imaging and spectroscopy data are based on observations with the NASA/ESA *Hubble Space Telescope*, obtained at the Space Telescope Science Institute, which is operated by AURA Inc., under NASA contract NAS 5-26555. This work was supported by grant GO-09793.01-A, GO-09793.03-A and GO-09793.08-A from the Space Telescope Science Institute. This project has made use of the aXe extraction software, produced by ST-ECF, Garching, Germany.

## REFERENCES

- Abraham, R.G., Ellis, R.S., Fabian, A.C. et al., 1999, *MNRAS*, 303, 641
- Abraham, R.G., van den Bergh, S., Glazebrook, K. et al., 1996, *ApJS*, 107, 1
- Alexander, D.M., Bauer, F.E., Brandt, W.N. et al., 2003, *AJ*, 126, 539
- Barnes, J.E., 1992, *ApJ*, 393, 484
- Beckwith, S.V.W. et al., 2005, in preparation
- Bell, E.F., Wolf, C., Meisenheimer, K. et al., 2004, *ApJ*, 608, 752
- Bender, R., Döbereiner, S., Möllenhoff, C., 1988, *A&A Suppl. Ser.*, 74, 385
- Bender, R., Surma, P., Döbereiner, S., Möllenhoff, C., Madejsky, R., 1989, *A&A*, 217, 35
- Bernardi, M., Sheth, R.K., James, A. et al., 2003, *AJ*, 125, 1882
- Bershady, M.A., Jangren, J.A., Conselice, C.J., 2000, *AJ*, 119, 2645
- Bertin, E. & Arnouts, S., 1996, *A&AS*, 117, 393
- Bertoldi, F., Carilli, C.L., Menten, K.M. et al., 2000, *A&A*, 360, 92
- Bolzonella, M., Miralles, J.-M., Pelló, R., 2000, *A&A*, 363, 476
- Bower, R.G., Lucey, J.R., Ellis, R.S., 1992, *MNRAS*, 254, 601
- Brichmann, J., Ellis, R.S., 2000, *ApJ*, 536, L77
- Bruzual, G. & Charlot, S., 2003, *MNRAS*, 344, 1000
- Caon, N., Capaccioli, M., D'Onofrio, M., 1993, *MNRAS*, 265, 1013
- Cappellari, M., Copin, Y., 2003, *MNRAS*, 342, 345
- Cimatti, A., Mignoli, M., Daddi, E. et al., 2002, *A&A*, 392, 395
- Chapman, S.C., Blain, A.W., Ivison, R.J., Smail, I., 2003, *Nature*, 422, 695
- Chapman, S.C., Windhorst, R.A., Odewahn, S., Yan, H., Conselice, C.J., 2004, *ApJ*, 599, 92
- Chen, H.-W., Marzke, R.O., McCarthy, P.J. et al., 2003, *ApJ*, 586, 745
- Colley, W.N., Rhoads, J.E., Ostriker, J.P., Spergel, D.N., 1996, *ApJL*, 473, L63
- Conselice, C.J., 2003, *ApJS*, 147, 1
- Conselice, C.J., Bershady, M.A., Dickinson, M., Papovich, C., 2003, *AJ*, 126, 1183
- Conselice, C.J., Bershady, M.A., Jangren, J.A., 2000, *ApJ*, 529, 886
- Conselice, C.J., Blackburne, J.A., Papovich, C., 2005, *ApJ*, 620, 564
- Cross, N.J., Bouwens, R.J., Benítez, N. et al., 2004, *AJ*, 128, 1990
- Daddi, E., Renzini, A., Pirzkal, N. et al., 2005, *ApJ*, 626, 680
- Dey, A., van Breugel, W.J.M., Vacca, W., Antonucci, R., 1997, *ApJ*, 490, 698
- Di Tullio, G.A., 1978, *A&A*, 62, L17
- Di Tullio, G.A., 1979, *A&A Suppl. Ser.*, 37, 591
- Dressler, A., 1980, 236, 351
- Dressler, A., Oemler, Ag.Jr., Couch, W.J. et al., 1997, *ApJ*, 490, 577
- Driver, S.P., Fernandez-Soto, A., Couch, W.J. et al., 1998, *ApJ*, 496, 93
- Driver, S.P., Windhorst, R.A., Griffiths, R., 1995, *ApJ*, 453, 48
- Ebner, K., Balick, B., 1985, *AJ*, 90, 183
- Eggen, O.J., Lynden-Bell, D., Sandage, A.R., 1962, *ApJ*, 136, 748
- Fasano, G., Cristiani, S., Arnouts, S. & Filippi, M., 1998, *AJ*, 115, 1400
- Fassnacht, C.D., Moustakas, L.A., Casertano, S. et al., 2004, *ApJ*, 600, L155
- Ferreras, I., Lisker, T., Carollo, C.M., Lilly, S.J., Mobasher, B., 2005, *astro-ph/0504127*
- Ferreras, I. & Silk, J., 2000, *MNRAS*, 316, 786
- Franx, M., 1988, *MNRAS*, 231, 285
- Galletta, G., 1980, *A&A*, 81, 179
- Genzel, R., Baker, A.J., Tacconi, L.J. et al., 2003, *ApJ*, 584, 633
- Giacconi, R., Zirm, A., Wang, J.X. et al., 2002, *ApJS*, 139, 369
- Giavalisco, M., Ferguson, H.C., Koekemoer, A.M. et al., 2004, *ApJ*, 600, 93
- Gilli, R., Cimatti, A., Daddi, E. et al., 2003, *ApJ*, 592, 721
- Glazebrook, K., Ellis, R., Santiago, B., Griffiths, R., 1995, *MNRAS*, 275, L19
- Goto, T., 2005, *MNRAS*, 357, 937
- Graham, A., Erwin, P., Caon, N., Trujillo, I., 2001, *ApJ*, 563, L11
- Graham, A., Lauer, T.R., Colless, M., Postman, M., 1996, *ApJ*, 465, 534
- Hamilton, D., 1985, *ApJ*, 297, 371
- Hernquist, L., 1992, *ApJ*, 400, 460
- Jedrzejewski, R.I., 1987, *MNRAS*, 226, 747
- Jesseit, R., Naab, Y., Burkert, A., 2005, submitted to *MNRAS*, *astro-ph/0501418*
- Jørgensen, I., Franx, M. & Kjørgaard, P., 1995, *MNRAS*, 273, 1097
- Kauffmann, G. & Charlot, S., 1998, *MNRAS*, 297, L23
- Kauffmann, G., White, S.D.M., Guiderdoni, B., 1993, *MNRAS*, 264, 201
- Kelson, D.D., Illingworth, G.D., van Dokkum, P.G., Franx, M., 2000, *ApJ*, 531, 184
- Khochfar, S., Burkert, A., 2005, *MNRAS*, 359, 1379
- Kormendy, J., 1977, *ApJ*, 218, 333
- Lauer, T.R., 1985, *MNRAS*, 216, 429
- Lauer, T.R., Faber, S.M., Gebhardt, K. et al., 2005, *AJ*, 129, 2138
- Le Fèvre, O., Guzzo, L., Meneux, B. et al., 2004, *A&A*, *astro-ph/0403628*
- Malin, D.F., Carter, D., 1983, *ApJ*, 274, 534
- Martel, A.R., Ford, H.C., Bradley, L.D. et al., 2004, *AJ*, 128, 2758
- Menanteau, F., Abraham, R. G. & Ellis, R. S., 2001, *MNRAS*, 322, 1
- Menanteau, F., Martel, A.R., Tozzi, P. et al., 2005, *ApJ*, 620, 697
- Michard, R., Poulain, P., 2000, *A&AS*, 141, 1
- Michard, R., Prugniel, P., 2004, *A&A*, 423, 833
- Mobasher, B., Idzi, R., Benítez, N. et al., 2004, *ApJ*, 600, L167
- Moustakas, L.A., Casertano, S., Conselice, C.J. et al., 2004, *ApJ*, 600, 131
- Naab, T., Burkert, A., 2003, *ApJ*, 597, 893
- Naab, T., Burkert, A. & Hernquist, L., 1999, *ApJ*, 523, L133
- Nelan, J.E., Smith, R.J., Hudson, M.J. et al., 2005, *ApJ*, accepted (*astro-ph/0505301*)
- O'Dea, C.P., Koekemoer, A.M., Baum, S.A. et al., 2001, *ApJ*, 121, 1915
- Odewahn, S.C., Burstein, D., Windhorst, R.A., 1997, *AJ*, 114, 2219
- Papovich, C., Dickinson, M., Ferguson, H.C., 2001, *ApJ*, 559, 620
- Papovich, C., Giavalisco, M., Dickinson, M. et al., 2003, *ApJ*, 598, 827
- Peletier, R.F., Balcells, M., 1996, *AJ*, 111, 2238
- Pettini, M., Shapley, A.E., Steidel, C.C. et al., 2001, *ApJ*, 554, 981
- Pirzkal, N., Xu, C., Malhotra, S. et al., 2004, *ApJS*, 154, 501
- Quinn, P.J., 1984, *ApJ*, 279, 596
- Rest, A., van den Bosch, F.C., Jaffe, W. et al., 2001, *AJ*, 121, 2431
- Sadler, E.M., Gerhard, O.E., 1985, *MNRAS*, 214, 177
- Sèrsic, J.-L., 1968, *Atlas de Galaxies Australes* (Cordoba: Obs. Astron.)
- Schade, D., Lilly, S.J., Crampton, D. et al., 1999, *ApJ*, 525, 31
- Shapley, A.E., Steidel, C.C., Adelberger, K.L., Dickinson, M., Giavalisco, M., Pettini, M., 2001, *ApJ*, 562, 95
- Smail, I., Ivison, R.J., Blain, A.W., Kneib, J.-P., 2002, *MNRAS*, 331, 495

- Stanford, S. A., Eisenhardt, P. R. & Dickinson, M. 1998, *ApJ*, 492, 461
- Steidel, C.C., Giavalisco, M., Pettini, M., Dickinson, M., Adelberger, K.L., 1996, *ApJ*, 462, L17
- Thielemann, F.-K., Nomoto, K. & Hashimoto, M., 1996, *ApJ*, 460, 408
- Thomas, D., 1999, *MNRAS*, 306, 655
- Thomas, D., Greggio, L. & Bender, R., 1999, *MNRAS*, 302, 537
- Thomas, D., Maraston, C., Bender, R., de Oliveira Mendes, C., 2005, *ApJ*, 621, 673
- Toomre, A., Toomre, J., 1972, *ApJ*, 178, 623
- Trager, S.C., Faber, S.M., Worthey, G., González, J.J., 2000, *AJ*, 119, 1645
- Tran, H.D., Tsvetanov, Z., Ford, H.C. et al., 2001, *AJ*, 121, 2928
- Treu, T., Ellis, R.S., Liao, T.X. et al., 2005, *ApJ*, submitted, astro-ph/0503164
- van Breugel, W.J.M., Stanford, S.A., Spinrad, H., Stern, D., Graham, J.R., 1998, *ApJ*, 502, 614
- van den Bosch, F.C., Ferrarese, L., Jaffe, W., Ford, H.C., O'Connell, R.W., 1994, *AJ*, 108, 1579
- van den Hoek, L. B. & Groenewegen, M. A. T., 1997, *A&AS*, 123, 305
- van Dokkum, P.G., Franx, M., 1995, *AJ*, 110, 2027
- van Dokkum, P.G., Franx, M., 2001, *ApJ*, 553, 90
- van Dokkum, P.G., Franx, M., Kelson, D.D., Illingworth, G.D., 1998, *ApJ*, 504, L17
- Vanzella, E., Cristiani, S., Dickinson, M. et al., 2005, *A&A*, 434, 53
- Yang, Y., Zabludoff, A.I., Zaritsky, D., Lauer, T.R., Mihos, J.C., 2004, *ApJ*, 607, 258
- Waddington, I., Windhorst, R.A., Cohen, S.H. et al., 2002, *MNRAS*, 336, 1342
- Webb, T.M.A., Eales, S.A., Lilly, S.J. et al., 2003, *ApJ*, 587, 41
- Williams, R.E., Blacker, B., Dickinson, M. et al., 1996, *AJ*, 112, 1335
- Windhorst, R.A., Taylor, V.A., Jansen, R.A. et al., 2002, *ApJS*, 143, 113
- Worthey, G., 1994, *ApJS*, 95, 107

TABLE 1. THE SAMPLE: UDF/GOODS IDS, COORDINATES AND MAGNITUDES (AB)

| ID                  | RA(J2000)  | Dec(J2000)  | F435W | F606W | F775W | F850LP | $z_{GRISM}$ | $z_{VLT}$ |
|---------------------|------------|-------------|-------|-------|-------|--------|-------------|-----------|
| J033229.22–274707.6 | 03:32:29.2 | –27:47:07.6 | 22.22 | 22.86 | 20.97 | 20.99  | 0.67        | 0.67      |
| UDF 2387            | 03:32:35.8 | –27:47:58.8 | 24.52 | 22.13 | 20.74 | 20.29  | 0.66        | 0.66      |
| UDF 1960            | 03:32:36.0 | –27:48:11.9 | 23.95 | 22.55 | 21.59 | 21.29  | 0.60        | 0.60      |
| UDF 9264            | 03:32:37.2 | –27:46:08.1 | 22.78 | 23.46 | 21.96 | 20.89  | 1.07        | 1.10      |
| UDF 3677            | 03:32:37.3 | –27:47:29.3 | 24.21 | 21.74 | 20.30 | 19.81  | 0.65        | 0.67      |
| UDF 8316            | 03:32:38.4 | –27:46:31.9 | 22.33 | 23.48 | 22.21 | 21.78  | 0.62        |           |
| UDF 6747            | 03:32:38.8 | –27:46:48.9 | 25.22 | 23.03 | 21.73 | 21.28  | 0.54        | 0.62      |
| UDF 68              | 03:32:38.8 | –27:49:28.5 | 26.83 | 24.49 | 22.98 | 22.36  | 0.82        |           |
| UDF 2322            | 03:32:39.2 | –27:47:58.4 | 24.93 | 22.49 | 21.07 | 20.62  | 0.66        |           |
| UDF 8               | 03:32:39.5 | –27:49:28.3 | 24.85 | 22.68 | 21.39 | 20.96  | 0.65        |           |
| UDF 6027            | 03:32:39.6 | –27:47:09.1 | 26.19 | 24.96 | 23.71 | 22.79  | 1.15        | 1.32      |
| UDF 153             | 03:32:39.6 | –27:49:09.6 | 25.15 | 22.95 | 21.51 | 20.63  | 0.98        | 0.98      |
| J033240.33–274957.0 | 03:32:40.3 | –27:49:57.0 |       | 23.51 | 22.23 | 21.88  | 0.66        |           |
| UDF 4587            | 03:32:40.7 | –27:47:31.0 | 23.99 | 22.96 | 21.67 | 21.22  | 0.67        | 0.67      |
| UDF 4527            | 03:32:41.4 | –27:47:17.2 | 24.42 | 22.41 | 21.06 | 20.61  | 0.62        |           |
| J033244.09–274541.5 | 03:32:44.1 | –27:45:41.5 | 23.11 | 21.12 | 20.09 | 19.75  | 0.49        |           |
| UDF 2107            | 03:32:45.8 | –27:48:12.9 | 24.42 | 22.77 | 21.76 | 21.38  | 0.62        | 0.53      |
| UDF 5177            | 03:32:48.5 | –27:47:19.6 | 24.74 | 23.33 | 22.43 | 22.08  | 0.50        |           |

TABLE 2. THE ISOPHOTAL PARAMETERS OF THE *bona fide* EARLY-TYPE GALAXIES. THESE HAVE NOT BEEN CORRECTED FOR THE ERRORS DERIVED IN THE APPENDIX.

| ID                  | Ellipticity | $\langle a_3/a \rangle \times 100$ | $\langle a_4/a \rangle \times 100$ | $\Delta PA$ |
|---------------------|-------------|------------------------------------|------------------------------------|-------------|
| J033229.22–274707.6 | 0.14        | –0.48                              | 0.23                               | 1.39        |
| UDF 2387            | 0.14        | –1.13                              | 0.83                               | 15.17       |
| UDF 1960            | 0.20        | 5.75                               | 3.59                               | 0.23        |
| UDF 9264            | 0.26        | –0.58                              | –0.59                              | 0.66        |
| UDF 3677            | 0.08        | –0.68                              | –0.65                              | 5.60        |
| UDF 8316            | 0.17        | 0.23                               | –0.15                              | 0.41        |
| UDF 6747            | 0.20        | 1.04                               | 3.61                               | 3.26        |
| UDF 68              |             |                                    |                                    |             |
| UDF 2322            | 0.13        | 0.65                               | 0.28                               | 1.96        |
| UDF 8               | 0.29        | –0.78                              | 0.47                               | 0.64        |
| UDF 6027            | 0.06        | 0.66                               | –0.77                              | 68.29       |
| UDF 153             | 0.26        | –0.10                              | –1.98                              | 15.90       |
| J033240.33–274957.0 |             |                                    |                                    |             |
| UDF 4587            | 0.41        | –1.93                              | 0.46                               | 0.50        |
| UDF 4527            | 0.36        | –0.11                              | 1.90                               | 6.69        |
| J033244.09–274541.5 | 0.30        | –0.48                              | 0.60                               | 4.76        |
| UDF 2107            | 0.22        | –0.50                              | 1.53                               | 0.07        |
| UDF 5177            | 0.19        | –0.85                              | 2.05                               | 37.28       |

## APPENDIX

## ERRORS ON THE MEASURED ISOPHOTAL PARAMETERS

When comparing the isophotal parameters of galaxies at different redshift and observed with different instruments, it becomes important to qualitatively establish the systematics in the data. For this reason, we simulated elliptical galaxies using a Sèrsic profile (Sèrsic 1968) with an index chosen between 3 and 4, an integrated magnitude and corresponding S/N ratio taken from the range spanned by the data. The input ellipticities and position angles were chosen among those measured for the *bona fide* early-type galaxies and the input  $a_3/a$  and  $a_4/a$  parameters were rigorously set to 0. The simulated galaxies were then convolved by the ACS PSF in the F775W filter; this PSF is the average image of stars in the UDF F775W frame.

We first ran ELLIPSE on the set of simulated galaxies convolved by the PSF and measured the isophotal parameters with the method described in Sect. 5. We then repeated the measurements on the simulated galaxies NOT convolved by the PSF. The results are summarized in Figures 12 and 13. When no PSF is applied, the output isophotal parameters agree extremely well with their respective input values, thus indicating that the S/N ratio of the data does not significantly affect our measurements (but see also Odewahn et al. 1997). The convolution with the PSF introduces deviations from the input parameters so that:

- i)* the output ellipticities are underestimated by about 10%;
- ii)* the output PA values are underestimated by about 3%;
- iii)* the output  $a_3/a$  parameter is scattered by about 0.2 around the input null value;
- iv)* the output  $a_4/a$  parameter is also overestimated by an amount which increases with the input ellipticity. For  $\epsilon < 0.5$  (the range in Table 2) we derive a discrepancy of +0.15 in  $a_4/a$ .

These deviations do not show any correlation with the input magnitude and Sèrsic index and represent the sistematic uncertainties on the values reported in Table 2.




FIG. 1.- Asymmetry and Concentration values for the *bona fide* early-type galaxies. The distinction between early-, late-type, mid galaxies and mergers is from Conselice et al. (2005).




FIG. 2.- Spectral energy distributions of the 18 galaxies in our sample (see Table 1. Each one is labeled with the UDF numbers, except for those outside of UDF, for which we give part of the GOODS identification. The error bars are the observed ACS/G800L data and the lines are the best fits according to the CSP models (see text for details).




FIG. 3.- Ages and metallicities corresponding to the best fit according to a simple exponentially decaying model (EXP; left) or a consistent chemical enrichment code (CSP; right). The dots are the average values of age or metallicity and the error bars represent the RMS of the distribution. Notice the EXP models assume a fixed metallicity for each star formation history. The solid lines in the bottom panel track the age of the Universe at a given redshift for a concordance cosmology. The dashed lines – from top to bottom – correspond to formation redshifts of  $z_F = \{5, 3, 2\}$ .




FIG. 4.- Rest-frame  $U - V$  vs  $M_V$  color-magnitude relation. The filled dots correspond to our sample after a K-correction to restframe  $U$  and  $V$  passbands. The hollow circles are the  $z = 0$  color-magnitude relation of the same sample. In order to evolve the stellar populations from their observed redshifts, we use the CSP model that gives the best fit. Typical error bars (including the effect of the K-correction) are shown. The triangles are Coma cluster galaxies from Bower et al. (1992) and the dashed lines delimit the best fit and scatter of this local cluster.




FIG. 5.-  $B$ -band Kormendy relation of our sample (black dots) compared to Coma cluster elliptical galaxies (squares; Jørgensen et al. 1995) and early-type galaxies from the HDF (grey dots; Fasano et al. 1998). The stars are the LBDS radio elliptical galaxies at  $z \sim 1.5$  from Waddington et al. (2002), while the open dots represent the early-type galaxies at  $z \sim 2$  studied by Daddi et al. (2005).




FIG. 6.- The isophotal parameters of the *bona fide* early-type galaxies (black, filled circles) and of the nearby ellipticals studied by Bender et al. (1988, 1989, grey and filled circles). These parameters are also plotted as a function of the galaxy half-light radius  $r_{hl}$  and total absolute magnitude in restframe B band  $M_B$ . The ellipticity and the  $a_4/a$  parameter of the *bona fide* early-type galaxies have been corrected for the PSF, i.e. ellipticities have been increased by 10% and  $a_4/a$  decreased by 0.15.

FIG. 7.- The (V-I) color maps of the *bona fide* early-type galaxies. Available at <http://www.stsci.edu/science/grapes/papers/pasquali05>.



FIG. 8.- As in Figure 7. Available at <http://www.stsci.edu/science/grapes/papers/pasquali05>.

FIG. 9.- *Top:* A close-up of the central region of J033244.09-274541.5 (left panel) and its extinction map (right panel) showing no presence of substructures. *Middle:* A close-up of the central region of UDF2387 and the corresponding extinction map, showing in white dust features in absorption. *Bottom:* A close-up of the central region of UDF3677 and its extinction map, with dust features in absorption. Close-ups are in units of counts/s in the F775W band, extinction maps are in units of magnitude. Available at <http://www.stsci.edu/science/grapes/papers/pasquali05>.

FIG. 10.- As in Figure 9, now showing substructures in emission. UDF4527 and UDF6747 are characterized by disk-like features, while UDF153 and UDF5177 represent galaxies with blue clumps and peculiar features, respectively. Available at <http://www.stsci.edu/science/grapes/papers/pasquali05>.




FIG. 11a.- The ELLIPSE parameters: surface brightness, ellipticity, PA,  $a_3/a$ ,  $a_4/a$  as a function of distance from the center of UDF6747 and derived in each of the F606W, F775W and F850LP filters. This galaxy has a remarkable disk substructure (cf. Figure 10). Figures 11b - 11r are available in the electronic edition of the Journal. The printed edition contains only Figure 11a as an example.




FIG. 12.- The comparison between the input and output ellipticities and PAs, for the simulated galaxies with (right-hand side panels) and without (left-hand side panels) convolution with the ACS PSF in the F775W.




FIG. 13.- The comparison between the input and output  $\langle a_3/a \rangle$  and  $\langle a_4/a \rangle$  for the simulated galaxies with (right-hand side panels) and without (left-hand side panels) convolution with the ACS PSF in the F775W.